




American Society of Hematology

Helping hematologists conquer blood diseases worldwide



The Menin Inhibitor Revumenib (SNDX-5613) Leads to Durable Responses in Patients with *KMT2A*-Rearranged or *NPM1* Mutant AML: Updated Results of a Phase 1 Study



Ghayas C. Issa, MD,¹ Ibrahim Aldoss, MD,² John F. DiPersio, MD, PhD,³ Branko Cuglievan, MD,¹ Richard M. Stone, MD,⁴ Martha L. Arellano, MD,⁵ Michael Thirman, MD,⁶ Manish R. Patel, MD,⁷ David Dickens, MD,⁸ Shalini Shenoy, MD,³ Neerav Shukla, MD,⁹ Galit Rosen, MD,¹⁰ Rebecca G. Bagley, MA,¹⁰ Michael L. Meyers, MD, PhD,¹⁰ Kate Madigan, MD,¹⁰ Peter Ordentlich, PhD,¹⁰ Yu Gu, PhD,¹⁰ Steven Smith, BS,¹⁰ Gerard M. McGeehan, PhD,¹⁰ and Eytan M. Stein, MD⁹

¹The University of Texas MD Anderson Cancer Center, Houston, TX; ²City of Hope, Duarte, CA; ³Washington University School of Medicine, St. Louis, MO; ⁴Dana-Farber Cancer Institute, Boston, MA; ⁵Emory University School of Medicine, Atlanta, GA; ⁶University of Chicago, Chicago, IL; ⁷Florida Cancer Specialists/Sarah Cannon Research Institute, Sarasota, FL; ⁸University of Iowa, Iowa City, IA; ⁹Memorial Sloan Kettering Cancer Center, New York, NY; ¹⁰Syndax Pharmaceuticals, Inc., Waltham, MA

Revumenib (SNDX-5613) is a potent, selective menin-KMT2A interaction inhibitor

- The menin-KMT2A interaction is a critical dependency in *KMT2Ar* (*MLL1r*) and *mNPM1* leukemias responsible for the leukemogenic gene expression
 - *KMT2Ar*: ~ 10% AML or ALL (~ 80% infant ALL)¹
 - *mNPM1*: ~ 30% AML²
- Revumenib competitively binds a discrete, well-defined pocket within menin, where both wild-type KMT2A (*MLL1*) and KMT2A fusion proteins bind

→ disassembling abnormal transcription complexes in *KMT2Ar*, *mNPM1*, and other leukemia subtypes³




1. Issa GC, et al. *Leukemia*. 2021;35:2482–2495; 2. Papaemmanuil, E. et al. *N Engl J Med*. 2016;374: 2209–2221; 3. Krivtsov A, et al. *Cancer Cell*. 2019;36(6):660–673.

ALL, acute lymphoblastic leukemia; AML, acute myeloid leukemia; *KMT2Ar*, lysine methyltransferase 2A rearrangements; *MLLr*, mixed lineage leukemia rearranged; *mNPM1*, mutated nucleophosmin 1.

AUGMENT-101 Phase 1 study design and objectives

Treatment

- Revumenib oral, q12h continuous dosing, 28-day cycle
- Accelerated titration into a rolling 6 design



Primary objectives

- Characterize safety, tolerability, PK, and RP2D

Prespecified RP2D selection criteria

- No more than 1 of 6 of evaluable patients experience a DLT
- At least 2/3 of patients receive $\geq 80\%$ of their dose in the first two cycles
- 24-hour AUC (AUC₀₋₂₄) exceeds 15,000 ng $\times$ hr/mL in at least 2/3 of patients

Exploratory objectives

- Anti-leukemic activity within *KMT2Ar* or *mNPM1* population

Populations

- Safety (pts ≥ 1 dose revumenib): N=68
- Efficacy (*KMT2Ar* or *mNPM1*): n=60

*Protocol originally allowed any R/R leukemia regardless of genotype but was amended to *KMT2Ar* or *mNPM1* patients only.

A majority of patients (n=60; 88%) were *KMT2Ar* or *mNPM1* and evaluable as the efficacy population.

AUC, area under the curve; DLT, dose-limiting toxicity; PK, pharmacokinetics; q12h, every 12 hours; RP2D, recommended Phase 2 dose; R/R, relapsed or refractory.

AUGMENT-101 patients are heavily pretreated with a poor prognosis

Baseline Characteristics	Safety Population N=68
Median age, years (range)	42.5 (0.8, 79)
Adult (n=60)	50.5
Pediatric (n=8)	2.5
Female, n (%)	42 (62)
Leukemia type, n (%)	
AML	56 (82)
ALL	11 (16)
MPAL	1 (2)
Median prior therapies (range)	4 (1,12)
Stem cell transplant, n (%)	31 (46)
Venetoclax, n (%)	41 (60)

Baseline Characteristics	Safety Population N=68
<i>KMT2A</i>Ar, n (%)	46 (68)
t(9;11)	10 (15)
t(11;19)	9 (13)
t(4;11)	6 (9)
t(6;11)	3 (4)
t(11;17)	2 (3)
Other	16 (24)
<i>mNPM1</i>, n (%)	14 (21)
<i>KMT2A</i> and <i>NPM1</i> wild type, n (%)	8 (12)
Co-occurring mutations*, n (%)	
<i>FLT3</i>	14 (25)
<i>RAS</i>	12 (29)
<i>TP53</i>	4 (10)

*In patients for whom co-occurring mutation data were available.
MPAL, mixed-phenotype acute leukemia

Data cutoff: 31 March 2022



2 patients with treatment ongoing and 12 patients proceeded to HSCT while in remission

Patient Disposition	Safety Population N=68
Treatment Ongoing, n (%)	2 (3)
Discontinued Treatment, n (%)	66 (97)
Progressive disease/No response	39 (57)
Transplant	12 (18)
Adverse event (all unrelated)	7 (10)
Withdrew consent	3 (4)
Other*	3 (4)
Physician decision	2 (3)

AUGMENT-101: Post-transplant outcomes
Abstract #376, Issa GC. et al.
Session 723; December 10, 4:00-5:30p

*Other: death (not related to treatment), n=2; donor lymphocyte infusion, n=1
HSCT, hematopoietic stem cell transplant.

Data cutoff: 31 March 2022



Adverse Events across all doses of revumenib

Any-grade treatment-related AE ($\geq 5\%$)	Safety Population N=68
Patients with ≥ 1 treatment-related AE, n (%)	53 (78)
ECG QTc prolonged	36 (53)
Nausea	18 (27)
Vomiting	11 (16)
Differentiation syndrome	11 (16)
Diarrhea	7 (10)
Dysgeusia	5 (7)
Decreased appetite	5 (7)

No treatment discontinuations for QTc prolongations, or associated arrhythmias

$\geq$ Grade 3 treatment-related AE	Safety Population N=68
Patients with $\geq$ Gr 3 treatment-related AE, n (%)	11 (16)
ECG QTc prolonged	9 (13)
Diarrhea	2 (3)
Anemia	2 (3)
Asthenia	1 (2)
Fatigue	1 (2)
Hypercalcemia	1 (2)
Hypokalemia	1 (2)
Neutropenia	1 (2)
Thrombocytopenia	1 (2)
Tumor lysis syndrome	1 (2)

10% of patients (5/52) had Gr 3 QTc prolongation at doses meeting criteria for RP2D

ECG, electrocardiogram; QTc, corrected QT interval.

Data cutoff: 31 March 2022

Revumenib demonstrates promising antileukemic activity in relapsed/refractory *KMT2Ar* and *mNPM1* leukemias

Best Response, n (%)	Efficacy Population n=60	
ORR[*]	32/60 (53%)	
Best Response		
CR	12 (20%)	CR/CRh 18 (30%)
CRh	6 (10%)	
CRp	5 (8%)	
MLFS	9 (15%)	
MRD^{neg} rate[†]	18/32 (56%)	
CR/CRh MRD ^{neg}	14/18 (78%)	
CR/CRh/CRp MRD ^{neg}	18/23 (78%)	
Genetic alteration	<i>KMT2Ar</i> n=46	<i>mNPM1</i> n=14
ORR	27/46 (59%)	5/14 (36%)
CR/CRh	15 (33%)	3 (21%)
CR/CRh MRD ^{neg} rate	11/15 (73%)	3/3 (100%)

^{*}Overall Response Rate = CR + CRh + CRp + MLFS; [†]MRD status assessed locally by PCR or MCF

CR, complete remission; CRh, complete remission with partial hematologic recovery; CRp, complete remission with incomplete platelet recovery; MLFS, morphologic leukemia free state; MRD, measurable residual disease.

Data cutoff:
31 March 2022

Revumenib demonstrates promising antileukemic activity in relapsed/refractory *KMT2Ar* and *mNPM1* leukemias

Best Response, n (%)	Efficacy Population n=60		Efficacy Population Doses Meeting Criteria for RP2D n=48	
ORR*	32/60 (53%)		25/48 (52%)	
Best Response				
CR	12 (20%)		8 (17%)	
CRh	6 (10%)		5 (10%)	
CRp	5 (8%)		5 (10%)	
MLFS	9 (15%)		7 (15%)	
MRD^{neg} rate[†]	18/32 (56%)		14/25 (56%)	
CR/CRh MRD ^{neg}	14/18 (78%)		10/13 (77%)	
CR/CRh/CRp MRD ^{neg}	18/23 (78%)		14/18 (78%)	
Genetic alteration	<i>KMT2Ar</i> n=46	<i>mNPM1</i> n=14	<i>KMT2Ar</i> n=37	<i>mNPM1</i> n=11
ORR	27/46 (59%)	5/14 (36%)	20/37 (54%)	5/11 (46%)
CR/CRh	15 (33%)	3 (21%)	10 (27%)	3 (27%)
CR/CRh MRD ^{neg} rate	11/15 (73%)	3/3 (100%)	7/10 (70%)	3/3 (100%)


Data cutoff:
31 March 2022

*Overall Response Rate = CR + CRh + CRp + MLFS; [†]MRD status assessed locally by PCR or MCF

CR, complete remission; CRh, complete remission with partial hematologic recovery; CRp, complete remission with incomplete platelet recovery; MLFS, morphologic leukemia free state; MRD, measurable residual disease.




Duration of revumenib therapy in patients with *KMT2Ar* or *mNPM1*



*Other reasons for treatment discontinuation included no response, relapse, death, and donor lymphocyte infusion.

Data cutoff: 31 March 2022


Duration of CR/CRh response with revumenib treatment



DOR, duration of response; NR, not reached.

Data cutoff: 31 March 2022


Overall survival in revumenib treated patients with *KMT2Ar* or *mNPM1*



Data cutoff: 31 March 2022

Differentiation syndrome managed with hydroxyurea and steroids


- 71-year-old with *KMT2Ar* AML relapsed after an allogeneic stem cell transplant
- Received revumenib at 339 mg PO q12h (Arm A), and achieved CRh, MRD negative remission



ANC, absolute neutrophil count; PB, peripheral blood; PO, by mouth; WBC, white blood cell.

Revumenib downregulates transcription of *FLT3* and induces responses in patients with *FLT3* co-mutations

- Among *KMT2Ar/mNPM1*: 14 had *mFLT3*
 - 13/14 (93%) previously treated with a *FLT3* inhibitor
- Overall response rate:
 - *KMT2Ar/mFLT3*: 2/3 (66%)
 - *mNPM1/mFLT3*: 3/11 (27%)
 - All responders previously treated with *FLT3* inhibitor
- Revumenib leads to transcriptional downregulation of *FLT3*, a putative target of *MEIS1*^{1,2}




- Responders
- Non-responders


Figure includes a subset of patients (WT and mutated) that were evaluable by gene expression at screening and C2D1.

1. Kühn MW, et al. Cancer Discov. 2016 Oct;6(10):1166-1181;
 2. Dzama MM, et al. Blood. 2020 Nov 19;136(21):2442-2456
 C, cycle; D, day; WT, wild type.

Revumenib response observed in extra-medullary disease



PET scan



CT scans of target lesions and spleen from a 19-year-old with relapsed *KMT2Ar* AML


- 3 prior lines of therapy, including 2 allogeneic stem cell transplants and local radiation to the spleen and abdominal nodes
- Received revumenib at 276 mg PO q12h
- Achieved CRh, MRD-negative remission with resolution of extramedullary disease in abdominal nodes and spleen

AP, anterior-posterior; FDG, fluorodeoxyglucose; PET, positron emission tomography; SUV, standardized uptake value.


Conclusions

- Revumenib resulted in deep, durable responses in heavily pre-treated R/R *KMT2Ar* and *mNPM1* patients, and demonstrated a clinically manageable safety profile
- 30% of patients attained CR/CRh with a median duration of 9.1 months
 - 78% of patients with CR/CRh attained MRD negativity
- 38% of responders proceeded to transplant
- Median OS was 7 months in this R/R population
- The only DLT, and the only common ($\geq 5\%$) $\geq$ Grade 3 related TEAE, was asymptomatic Grade 3 QTc prolongation
 - 10% in patients treated at doses meeting criteria for RP2D; 13% in patients treated at all doses tested
- Differentiation syndrome occurred in 16% of patients
 - All cases were Grade 2 and responded to management with steroids with or without hydroxyurea

AUGMENT-101 Phase 2 pivotal trials underway in 3 distinct patient populations



Dose:
Revumenib 163 mg q12h
with a strong CYP3A4 inhibitor



*Patients taken to HSCT can restart treatment with revumenib post-transplant.

Acknowledgements

We thank all AUGMENT-101 study patients, their families, and caregivers for participating in this trial.

Principal Investigators, staff, and sites where the trial was conducted:

Memorial Sloan Kettering Cancer Center

Eytan Stein

Kait Tkachuk

Dana-Farber Cancer Institute

Richard Stone

Emilie Baxter

Emory University School of Medicine

Martha Arellano

Temitayo Obajuluwa

University of Chicago

Michael Thirman

Adriana Koziol

Florida Cancer Specialists

Manish Patel

Terri Peterson

The University of Texas MD Anderson Cancer Center

Ghayas Issa

Branko Cuglievan

Allison Pike

Washington University School of Medicine

John DiPersio

Jennifer Reed

City of Hope

Ibrahim Aldoss

Manjyot Nanhwan

Montefiore Medical Center

Ioannis Mantzaris

David Loeb

Joel Victor

University of Iowa

David Dickens

Christopher Stamy

AUGMENT-101: Post-transplant outcomes

Abstract #376

Session 723. Allogeneic Transplantation: Long-term Follow-up and Disease Recurrence II

December 10, 4:00-5:30p

This research was sponsored by Syndax Pharmaceuticals, Inc.